



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

AVERAGE AND PROBABILITY.

113. Proposed by F. P. MATZ, Sc. D., Ph. D., Professor of Mathematics and Astronomy, Defiance College, Defiance, Ohio.

A given cube is cut by a plane in such a manner that the lines of section form a regular hexagon. What is the mean area of this hexagon?

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics in The Temple College, Philadelphia, Pa.

Let $ABCD-H$ be the cube, side a .

$HM=EN=AI=BJ=CK=GL=x$.

$LH=EM=AN=BI=CJ=GK=a-x$.

Then $IJKLMN$ is a hexagon.

$IJ=JK=KL=LM=MN=NI=\sqrt{(a^2+2x^2-2ax)}$.


$IK=JL=KM=LN=MI=NJ=\sqrt{(2a^2+2x^2-2ax)}$.

\therefore The hexagon is a regular hexagon.

Area of this hexagon $=\frac{3}{2}(a^2+2x^2-2ax)\sqrt{3}=u$.

Average area $=\Delta=\int_{\frac{1}{2}a}^a u dx / \int_{\frac{1}{2}a}^a dx$.

$\therefore \Delta = \frac{3\sqrt{3}}{a} \int_{\frac{1}{2}a}^a (a^2+2x^2-2ax) dx = a^2\sqrt{3}$.



114. Proposed by L. C. WALKER, A. M., Professor of Mathematics, Petaluma High School, Petaluma, Cal.

If a regular polygon of n sides be placed at random on another equal polygon, show that the chance that the center of the first will fall on the second polygon is

$$\frac{\pi}{2[\pi + n \tan(\pi/n)]}.$$

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics in The Temple College, Philadelphia, Pa.

Let AOB , DCE be a triangle of each polygon. Let DCE move parallel to itself, then the center D will trace out the polygon of $2n$ sides $KLMGN$, etc.

Let $BO=CD=r$, $\angle BCE=\theta$ =the angle BO makes with DC . Then the perpendicular distance from AB to $HG=r\cos(\pi/n)-\theta$, $HA=r\sec(\pi/n)\cos[(\pi/n)-\theta]=r[\cos\theta+\sin\theta\tan(\pi/n)]$, $OH=r[1+\cos\theta+\sin\theta\tan(\pi/n)]$.

$MG=\sqrt{(CD^2+BG^2-2CD\cdot BG\cos\theta)}=r\sin\theta\sec(\pi/n)$.

$HG=2r[1+\cos\theta+\sin\theta\tan(\pi/n)]\sin(\pi/n)$.

$HL=HG-LG=r[2\cos\theta\sin(\pi/n)+2\sin\theta\sin(\pi/n)\tan(\pi/n)-\sin\theta\sec(\pi/n)]$.

Area $HOG=\frac{1}{2}r^2[1+\cos\theta+\sin\theta\tan(\pi/n)]^2\sin(2\pi/n)$.

Area $MGN=\frac{1}{2}r^2[2\sin\theta\cos\theta\tan(\pi/n)+\sin^2\theta\tan^2(\pi/n)-\sin^2\theta]\sin^2(\pi/n)$.

The total number of positions is $n(\triangle OGH-\triangle MGN)=nr^2\sin(2\pi/n)[1+\cos\theta+\sin\theta\tan(\pi/n)]=A$. The number of favorable positions is $n\triangle AOB=B$.

$\therefore B=\frac{1}{2}nr^2\sin(2\pi/n)$.

$\therefore p = \int_0^{\pi/n} B d\theta / \int_0^{\pi/n} A d\theta = \frac{1}{2} \int_0^{\pi/n} d\theta / \int_0^{\pi/n} [1+\cos\theta+\sin\theta\tan(\pi/n)] d\theta = \frac{\pi}{2[\pi+n\tan(\pi/n)]}$

COR. When $n=\infty$, $p=\frac{1}{4}$; when $n=3$, $p=\pi/[2(\pi+3\sqrt{3})]$; when $n=4$, $p=\pi/[2(\pi+4)]$; when $n=6$, $p=\pi/[2(\pi+2\sqrt{3})]$.

